



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

Modulation of cell surface GABA(B) receptors by desensitization, trafficking and regulated degradation

Benke, Dietmar ; Zemoura, Khaled ; Maier, Patrick J

Abstract: Inhibitory neurotransmission ensures normal brain function by counteracting and integrating excitatory activity. -Aminobutyric acid (GABA) is the main inhibitory neurotransmitter in the mammalian central nervous system, and mediates its effects via two classes of receptors: the GABA(A) and GABA(B) receptors. GABA(A) receptors are heteropentameric GABA-gated chloride channels and responsible for fast inhibitory neurotransmission. GABA(B) receptors are heterodimeric G protein coupled receptors (GPCR) that mediate slow and prolonged inhibitory transmission. The extent of inhibitory neurotransmission is determined by a variety of factors, such as the degree of transmitter release and changes in receptor activity by posttranslational modifications (e.g., phosphorylation), as well as by the number of receptors present in the plasma membrane available for signal transduction. The level of GABA(B) receptors at the cell surface critically depends on the residence time at the cell surface and finally the rates of endocytosis and degradation. In this review we focus primarily on recent advances in the understanding of trafficking mechanisms that determine the expression level of GABA(B) receptors in the plasma membrane, and thereby signaling strength.

DOI: <https://doi.org/10.4331/wjbc.v3.i4.61>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-63558>

Journal Article

Published Version

Originally published at:

Benke, Dietmar; Zemoura, Khaled; Maier, Patrick J (2012). Modulation of cell surface GABA(B) receptors by desensitization, trafficking and regulated degradation. *World Journal of Biological Chemistry*, 3(4):61-72.

DOI: <https://doi.org/10.4331/wjbc.v3.i4.61>

Modulation of cell surface GABA_B receptors by desensitization, trafficking and regulated degradation

Dietmar Benke, Khaled Zemoura, Patrick J Maier

Dietmar Benke, Khaled Zemoura, Patrick J Maier, Institute of Pharmacology and Toxicology, University of Zürich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

Author contributions: All authors contributed to the writing of this review.

Supported by The Swiss Science Foundation Grant, 31003A_121963

Correspondence to: Dietmar Benke, PhD, Institute of Pharmacology and Toxicology, University of Zürich, Winterthurerstrasse 190, 8057 Zurich, Switzerland. benke@pharma.uzh.ch

Telephone: +41-44-6355930 Fax: +41-44-6356874

Received: August 27, 2011 Revised: December 8, 2011

Accepted: December 15, 2011

Published online: April 26, 2012

© 2012 Baishideng. All rights reserved.

Key words: GABA_B receptors; Neuron; Trafficking; Endocytosis; Recycling; Degradation

Peer reviewers: Francisco Ciruela, Dr., Department de Patologia i Terapèutica Experimental, Pavelló de Govern, L'Hospitalet de Llobregat 08907, Spain; Sung H Kim, Professor, College of Oriental Medicine, Kyunghee University, Cancer Preventive Material Research Center, Seoul 130-701, South Korea; Joe B Blumer, Assistant Professor, Department of Pharmacology, Medical University of South Carolina, 114 Doughty St., STB331, Charleston, SC 29425, United States; Saobo Lei, Associate Professor, Department of Pharmacology, Physiology and Therapeutics, University of North Dakota, 501 N Columbia Road, Grand Forks, ND 58203, United States

Abstract

Inhibitory neurotransmission ensures normal brain function by counteracting and integrating excitatory activity. γ -Aminobutyric acid (GABA) is the main inhibitory neurotransmitter in the mammalian central nervous system, and mediates its effects *via* two classes of receptors: the GABA_A and GABA_B receptors. GABA_A receptors are heteropentameric GABA-gated chloride channels and responsible for fast inhibitory neurotransmission. GABA_B receptors are heterodimeric G protein coupled receptors (GPCR) that mediate slow and prolonged inhibitory transmission. The extent of inhibitory neurotransmission is determined by a variety of factors, such as the degree of transmitter release and changes in receptor activity by posttranslational modifications (e.g., phosphorylation), as well as by the number of receptors present in the plasma membrane available for signal transduction. The level of GABA_B receptors at the cell surface critically depends on the residence time at the cell surface and finally the rates of endocytosis and degradation. In this review we focus primarily on recent advances in the understanding of trafficking mechanisms that determine the expression level of GABA_B receptors in the plasma membrane, and thereby signaling strength.

Benke D, Zemoura K, Maier PJ. Modulation of cell surface GABA_B receptors by desensitization, trafficking and regulated degradation. *World J Biol Chem* 2012; 3(4): 61-72 Available from: URL: <http://www.wjgnet.com/1949-8454/full/v3/i4/61.htm> DOI: <http://dx.doi.org/10.4331/wjbc.v3.i4.61>

FUNCTIONS OF GABA_B RECEPTORS

Metabotropic GABA_B receptors are widely distributed throughout the central nervous system where they mediate slow, prolonged inhibition to control neuronal excitation, and contribute to synaptic plasticity^[1].

GABA_B receptors are present at pre- and postsynaptic sites of both inhibitory and excitatory neurons. Electron microscopy revealed that GABA_B receptors are located predominantly at areas close to neurotransmitter release sites and at peri- and extrasynaptic areas of spines and dendrites, but only rarely directly at active zones or postsynaptic densities^[2-7]. This location of GABA_B receptors implies that they are not directly activated by synaptically released GABA. One mechanism to activate GABA_B receptors requires intense neuronal activity, resulting in




Figure 1 Structural organization of GABA_B receptors. Functional GABA_B receptors are heterodimers composed of the two subunits GABA_{B1} and GABA_{B2}. Both subunits are heptahelical membrane proteins with a large extracellular located N-terminal domain containing a "Venus flytrap" module and a large intracellular C-terminal domain containing a coiled-coil protein-protein interaction module. GABA_{B1} and GABA_{B2} heterodimerize via their "Venus flytrap" and coiled-coiled domains. An endoplasmic reticulum (ER) retention/retrieval signal is present distal to the coiled-coil domain in GABA_{B1} and prevents ER exit of GABA_{B1} unless it is masked by heterodimerization with GABA_{B2}. The "Venus flytrap" module of GABA_{B1} constitutes the GABA binding site, whereas that of GABA_{B2} is inactive and not involved in ligand binding. Instead, the heptahelical domain of GABA_{B2} contains a binding site for allosteric modulators, which affects the affinity of ligands binding to the GABA site. Binding of GABA results in the recruitment and activation of G α i/o proteins via GABA_{B2}. The activated G α i/o subunit inhibits the adenylyl cyclase, resulting in lowered cAMP levels, while the G β γ dimer activates K⁺ channels and inhibits Ca²⁺ channels, leading in either case to neuronal inhibition. There exist two isoforms of GABA_{B1}, named GABA_{B1a} and GABA_{B1b}, which are generated by alternative promoter usage. They only differ by the additional presence of two so-called "sushi repeats" (protein-protein interaction modules) in the N-terminal domain of GABA_{B1a}. GABA: γ -Aminobutyric acid; ATP: Adenosine-5'-triphosphate; cAMP: 3'-5'-cyclic adenosine monophosphate.

a spill-over of synaptically released GABA^[8]. However, there are also other sources that may increase the ambient level of GABA, such as activity-dependent release of GABA from dendrites and glia cells^[9-11]. Recently, it has been shown that basal synaptic activity generates a sufficient concentration of ambient GABA to tonically induce a low level of presynaptic GABA_B receptor activation, which results in the control of transmitter release^[12].

Binding of GABA to the GABA_B receptor activates Gi/o-type G proteins^[13-18], which in turn modulate three major effector systems: adenylyl cyclases, voltage-sensitive Ca²⁺

channels and inwardly-rectifying K⁺ channels (Figure 1).

The α subunit of the activated G protein inhibits adenylyl cyclase activity, which decreases cellular 3'-5'-cyclic adenosine monophosphate (cAMP) levels and affects the activity of cAMP-dependent processes. Unfortunately, the contribution of GABA_B receptor-induced lowering of cAMP levels to physiological processes is poorly investigated. So far it has been shown that it retards synaptic vesicle recruitment during sustained activity, which reduces transmitter release^[19]. In addition, GABA_B receptor-mediated G α i/o effects may be important for long-term

adaptations involving regulation of protein kinase activity and gene transcription^[20,22].

However, the most well established GABA_B receptor actions are mediated *via* the $\beta\gamma$ dimer of the activated G protein. At presynaptic sites, voltage-sensitive P/Q- and N-type Ca²⁺ channels are the predominant effectors of GABA_B receptors^[23,27]. GABA_B receptor activated G $\beta\gamma$ inhibits Ca²⁺ channel activity by slowing their current activation kinetics^[28], which eventually results in reduced transmitter release. Postsynaptically, GABA_B receptor effects are mainly mediated by the family of G protein-gated inwardly rectifying K⁺ channels (GIRK1-4 also called Kir3.1-3.4)^[29,30]. G $\beta\gamma$ directly binds to GIRK channels^[31,32] and activates them^[33,34], resulting in an outward K⁺ current. This hyperpolarizes the membrane and consequently inhibits neuronal activity. However, there is no strict mechanistic segregation of pre- (Ca²⁺ channels) and postsynaptic (K⁺ channels) effector systems. There is accumulating evidence that GABA_B receptors also activate K⁺ channels at presynaptic sites, which assists inhibition of transmitter release^[35,37]. Conversely, there is also data for GABA_B receptor mediated inhibition of postsynaptic Ca²⁺ channels^[38-41]. This provides an additional mechanism for controlling the excitability of dendrites and spines. Thus, the current data is consistent with a complex pattern of regulating the activity of multiple G protein-gated inwardly rectifying K⁺ channels and voltage-sensitive Ca²⁺ channels, both at pre- and postsynaptic sites, resulting in the inhibition of neuronal activity.

To ensure efficient activation of the effector system, GABA_B receptors are localized in close proximity to their effector channels^[36,42] and may even constitute signaling complexes by physical interaction^[36,43].

MOLECULAR ORGANIZATION OF GABA_B RECEPTORS

Although the GABA_B receptor was discovered in 1980^[44], its molecular identity and characterization was delayed for almost 20 years until the first constituent of the receptor was cloned. This delay was due to the fact that all biochemical attempts to purify the receptor failed and expression cloning proved unsuccessful. The development of high-affinity antagonists eventually permitted the successful screening of expression libraries yielding two cDNAs derived from a single gene, GABA_{B1a} and GABA_{B1b}^[45]. GABA_{B1a} and GABA_{B1b} are generated by differential promoter usage^[46] and differ solely by the presence of an additional N-terminal sequence in GABA_{B1a} coding for two protein-protein interaction domains, so-called “sushi domains”. GABA_{B1a} and GABA_{B1b} show all the characteristics of class III G protein-coupled receptors (e.g., a very large extracellular domain, seven transmembrane-spanning (heptahelical) sequences and a large intracellular located C-terminal domain) (Figure 1). So far, no functional differences among GABA_B receptors containing GABA_{B1a} and GABA_{B1b} have been detected. The cloning of these first GABA_B receptor constituents

provided the basis for numerous research efforts analyzing the molecular characterization and function of GABA_B receptors. It soon became clear that functional GABA_B receptors are obligatory heterodimers composed of GABA_{B1} (either GABA_{B1a} or GABA_{B1b}) and a second heptahelical membrane protein named GABA_{B2}, sharing about 35% sequence identity with GABA_{B1}^[47-51]. Both subunits serve distinct functions within the heterodimeric receptor complex. GABA_{B1} contains the agonist and antagonist binding site in the large N-terminal extracellular domain, which is most likely arranged in a Venus flytrap-like structure^[52-54]. Association with GABA_{B2} is necessary to keep the GABA binding site in a high affinity state^[55,56]. On the other hand, GABA_{B2} contains a binding site for allosteric modulators, which is not however associated with the N-terminal Venus flytrap domain, but is located in the heptahelical domain^[57]. Binding of ligands to this site does not directly activate the GABA_B receptor but instead affects the affinity of orthosteric agonists and antagonists to GABA_{B1}^[58]. Finally, GABA_{B2} is responsible for G protein activation^[56,59-63] and plays an important role in cell surface trafficking of the heterodimerized receptor complex by masking an arginine-based endoplasmic reticulum (ER) retention/retrieval (RXR) signal present in the C-terminal domain of GABA_{B1}^[64-68].

THE ROLE OF DESENSITIZATION AND PHOSPHORYLATION ON THE AVAILABILITY OF FUNCTIONAL GABA_B RECEPTORS

Prolonged exposure of G protein coupled receptors (GPCR) to agonists generally leads to a complex series of events in order to attenuate or terminate signal transduction, protecting the cell from overstimulation. Signal transduction is often attenuated by desensitization of the receptors (i.e., abrogating signaling), although the agonist is still present^[69,70]. Desensitization of many GPCRs involves phosphorylation-dependent uncoupling of the receptor from the G proteins, followed by internalization of the receptor. Activated GPCRs are usually phosphorylated by G protein-coupled receptor kinase (GRKs) at serine and/or threonine residues residing in the carboxyl-terminal tail- or intracellular loop regions, which rapidly attenuates receptor responses. Phosphorylation leads to the recruitment of arrestins, which is thought to sterically prohibit signaling to G proteins and induces internalization of the receptor by linking it to components (clathrin, AP2 complex) of the endocytosis machinery^[69,70]. Internalized receptors are then either degraded in lysosomes or are dephosphorylated and subsequently recycled to the plasma membrane, where they are again available for signaling.

It is well known that prolonged activation of GABA_B receptors commonly leads to their desensitization. Recent studies suggest that there might be more than one mechanism for desensitization of GABA_B receptors^[71-75],




Figure 2 Mechanisms of GABA_B receptor desensitization. Three distinct mechanisms have been so far implicated in the desensitization of GABA_B receptors. In cerebellar granule cells, G protein receptor kinase (GRK) 4 and 5 associate with GABA_B receptors and induce desensitization of the receptors in a phosphorylation-independent manner. In cortical and hippocampal neurons, desensitization of the receptors involves the interaction of NEM-sensitive fusion protein (NSF) with GABA_{B1} and GABA_{B2}, which is thought to prime the receptor for phosphorylation by protein kinase C (PKC). Association of potassium channel tetramerization domain-containing (KCTD) proteins 12 and 12b with the C-terminus of GABA_{B2} appears to render the receptor complex competent for desensitization. GABA: γ -Aminobutyric acid.

which does not follow the classical desensitization pattern of GPCRs described above (Figure 2). Distinct desensitization mechanisms for GABA_B receptors may be operative in different neuronal populations.

A study conducted in mouse cerebellar granule cells showed that GRK4, which is mainly expressed in testes and cerebellum^[69,76,77], promotes agonist-induced desensitization of GABA_B receptors *via* direct association, but does not involve GABA_B receptor phosphorylation^[78]. These findings were confirmed by Kainaide *et al.*^[75], who demonstrated that the association of GABA_{B2} with GRK4 or GRK5, but not GRK2, -3 or -6, leads to agonist-induced receptor desensitization in *Xenopus* oocytes and baby hamster kidney cells. Interestingly, GRK4 and -5-mediated desensitization was partially suppressed by application of S(+)-ketamine, which leads to inhibition of the GABA_B receptors/GRK complex formation by an as yet unidentified mechanism^[79]. It is currently not understood how GRK4 and -5 mediate desensitization of GABA_B receptors. However, it might well be that the binding of GRK4 and -5 disrupts GABA_B receptor/G-protein interaction.

On the other hand, for cortical and hippocampal neurons, a phosphorylation-dependent desensitization mechanism of GABA_B receptors was reported^[74]. This mechanism is based on the direct interaction of NEM-sensitive fusion (NSF) protein with the C-terminal domains of GABA_{B1} and GABA_{B2}, which primes the receptor for recruitment of protein kinase C (PKC). The data indicate that the association of GABA_B receptors with NSF is a prerequisite for recruiting PKC to the receptor upon agonist activation. PKC phosphorylates the receptor leading to its desensitization, and induces dissociation of NSF from the receptors. The precise roles of NSF and PKC in this complex process remain to be determined. NSF might be required for unmasking phosphorylation sites of the receptor or involved in PKC activation. In addition, it is unclear whether NSF dissociates from the receptor before desensitization occurs or whether releasing NSF initiates recovery of the receptor from desensitization.

Another factor determining desensitization of GABA_B receptors was recently discovered by functional proteomics^[80]. Members of the potassium channel tetramer-

ization domain-containing (KCTD) protein family were found to interact as tetramers with the C-terminus of GABA_{B2}, generating high-molecular mass protein complexes. Depending on the co-expressed KCTD subtype, distinct parameters of GABA_B receptor function were affected, such as agonist potency, signaling onset or desensitization. Interestingly, only when GABA_B receptors were co-expressed with KCTD-12 or -12b desensitization of GABA_B receptors was observed, whereas in the presence of KCTD-8 or -16 the receptors displayed no desensitization^[80]. This finding may explain the observation that GABA_B receptor desensitization varies among different neuronal populations. One striking example is the ventral tegmental area (VTA). GABA_B receptors expressed in GABAergic neurons of the VTA display baclofen-induced, largely non-desensitizing, currents, whereas in dopaminergic neurons of the VTA baclofen elicited desensitizing currents^[81]. These findings suggest that the general ability of GABA_B receptors to desensitize may be determined by the associated KCTD subtype, which may then recruit distinct desensitization mechanisms depending on the neuronal population.

Interestingly, PKA-dependent phosphorylation appears to counteract desensitization of GABA_B receptors. Couve *et al.*^[71] showed that PKA exclusively phosphorylates serine 892 (S892) in the C-terminal domain of GABA_{B2}, resulting in reduced receptor desensitization. This effect on desensitization can be overcome by activation of the receptors, which results in inhibition of adenylyl cyclases, reduced cAMP levels and consequently diminished PKA activity and GABA_{B2}-S892 phosphorylation. The precise mechanism as to how PKA phosphorylation of GABA_{B2}-S892 affects desensitization of GABA_B receptors remains unclear. There is an indication that it stabilizes cell surface GABA_B receptors and thereby increases effector coupling^[71,82]. This is, however, unlikely because it is now well accepted that prolonged agonist exposure does not trigger increased internalization of cell surface receptors^[78,82-85]. However, GABA_{B2}-S892 phosphorylation provides a mechanism for regulating the extent of GABA_B receptor desensitization by the activity of G α_s -coupled GPCRs that enhance PKA activity.

Another kinase that is involved in regulating GABA_B receptor activity is the 5'AMP-dependent protein kinase (AMPK). AMPK directly binds to the C-terminus of GABA_{B1} and phosphorylates S917 and S783 in the C-terminal domains of GABA_{B1} and GABA_{B2}, respectively^[86]. Functional analysis revealed that phosphorylation of S783 resulted in a stabilization of baclofen-induced K⁺ currents^[86]. This effect has been shown to be of particular relevance in limiting neuronal cell death in experimental ischemia. Anoxic or ischemic conditions are associated with neuronal over-excitation, a decline in cellular adenosine-5'-triphosphate (ATP) and a rise in Ca²⁺ and AMP levels, which are all factors activating AMPK^[87,88]. Under such conditions, increased phosphorylation of GABA_{B2}-S783 was detected along with an over-expression of a GABA_{B2} mutant that cannot be phosphorylated at this

site associated with increased neuronal death^[86]. These findings support a mechanism in which AMPK functions as a metabolic sensor that detects severe cellular stress and phosphorylates, amongst others, GABA_B receptors. This is thought to result in enhanced GABA_B receptor signaling that counteracts over-excitation of the neuron and limits neuronal death.

REGULATION OF GABA_B RECEPTORS BY TRAFFICKING

The lifecycle of a plasma membrane protein like the GABA_B receptor starts with its synthesis at the rough ER where the nascent protein is co-translationally incorporated into the ER membrane. After folding, initial posttranslational processing, and assembly, the receptor is exported to the Golgi apparatus where it is further processed and finally transported *via* the trans-Golgi network to the plasma membrane. After a certain time span of function, the receptor is internalized and recycled back into the plasma membrane for another cycle of function, or is eventually degraded into lysosomes. To ensure a constant number of receptors in the plasma membrane for signaling, these trafficking events need to be precisely coordinated. On the other hand, regulation of each of the different trafficking steps permits adjusting the number of cell surface receptors, and thus signaling strength, according to the physiological requirements.

ER export of GABA_B receptors

Little is known about the early stages in the lifecycle of GABA_B receptors. So far it is clear that exit of heterodimeric GABA_B receptors from the ER is controlled by an arginine-based ER retention/retrieval signal (RXR) present in the C-terminal domain of GABA_{B1}^[64-66]. The mechanism that prevents cell surface trafficking of GABA_{B1} appears to involve the coat protein complex I (COP I), which plays a central role in the retrograde transport of proteins from the Golgi apparatus back to the ER^[89]. COP I binds to the ER retention/retrieval signal of GABA_{B1} and shuttles monomeric GABA_{B1} that reached the cis-Golgi apparatus back to the ER. Heterodimerization with GABA_{B2} masks the ER retention/retrieval signal and permits forward transport^[64-68]. In contrast to GABA_{B1}, monomeric GABA_{B2} can leave the ER and reach the cell surface. However, it is assumed that the GABA_{B2} expression level in the ER is a limiting factor for ER exit of the heterodimeric GABA_B receptors. This mechanism is thought to ensure that only correctly folded and assembled (i.e., functional) receptors are exported to the cell surface.

Endocytosis of GABA_B receptors

There are two principal mechanisms by which GPCRs are internalized from the plasma membrane, constitutive endocytosis and agonist-induced endocytosis. Constitutive endocytosis constantly removes receptors from the

cell surface, whereas agonist-induced endocytosis initiates removal of receptors from the plasma membrane upon activation of the receptors and ensures fast termination of signaling. It is now well established that GABA_B receptors undergo constitutive endocytosis, whereas the presence of agonist-induced internalization of the receptors is less clear.

Heterologously expressed, as well as neuronal, GABA_B receptors display fast constitutive internalization, as evidenced by distinct experimental approaches including immunofluorescence staining and microscopy, live cell imaging and cell surface biotinylation methods^[83,84,90-93]. Constitutive internalization of GABA_B receptors is a fast process, as shown by the rapid loss of labeled receptors from the cell surface, which reaches a plateau after 10-30 min (40% of labeled receptors remain at the cell surface), with rates of internalization of ranging from 2-10 min^[92-94]. GABA_B receptors internalize as heterodimers and are not dissociated into its subunits prior to endocytosis^[84,90,92,95]. The rate of internalization appears to be determined by GABA_{B2}. GABA_{B1}, which contains an inactivated ER retention signal so that it is exported to the plasma membrane, displays a considerably faster rate of internalization than the GABA_{B1,2} heterodimer^[92]. This is due to a dileucine motif within the coiled-coil domain of GABA_{B1}, which gets masked upon assembly with GABA_{B2}.

The data so far suggest that for endocytosis, GABA_B receptors are recruited to clathrin-coated pits and internalized in a dynamin-dependent manner^[83,90,95]. Clathrin-coated pits are composed of clathrin heavy and light chains that form a polymeric lattice and contain numerous adaptor and endocytic accessory proteins. For endocytosis, the cargo-loaded clathrin-coated pit invaginates and is eventually released from the plasma membrane in a GTP-dependent reaction mediated by dynamin^[96]. There is evidence based on colocalization and immunoprecipitation data that GABA_B receptors interact with the AP2 adaptor, which is one of the adaptor complexes that recruit membrane proteins to clathrin-coated pits^[83,84,90].

Colocalization studies with marker proteins for various endosomal compartments revealed that endocytosed GABA_B receptors first enter early endosomes and are then either sorted to Rab4 or Rab11-positive recycling endosomes, or to Rab7-positive late endosomes, and finally to lysosomes for degradation^[84,90,92,95,97,98].

In addition to the colocalization data, there is also functional evidence that endocytosed GABA_B receptors constitutively recycle back to the cell surface. Using immunofluorescence staining and tagged GABA_B receptors transfected into hippocampal neurons Vargas *et al.*^[90] showed that a significant fraction of endocytosed receptors recycle back to the cell surface. Quantitative cell surface biotinylation and immunofluorescence-based methods indicate that the vast majority of native GABA_B receptors in cortical neurons are rapidly recycled to the plasma membrane. After 15 min, about half of the internalized receptors have recycled back to the cell surface, and after 30 min

this has increased to the majority of the receptors^[84,94].

In summary, the current data indicate that GABA_B receptors constitutively internalize at a high rate *via* the classical clathrin-dependent pathway and rapidly recycle back to the cell surface. Since endocytosis and recycling are highly energy-consuming processes, this mechanism is most likely of significant physiological relevance. The most obvious explanation is that a high rate of constitutive internalization and recycling generates a pool of intracellular receptors that can be immediately inserted into the plasma membrane to increase the cell surface number of receptors by increasing the rate of recycling while leaving the rate of internalization constant. In the case of synaptic AMPA receptors, such a mechanism has been proposed to contribute to increasing the level of the receptors during the early phase of long-term potentiation, which is thought to underlie learning and memory formation^[99].

Degradation of GABA_B receptors

Most cell surface receptors are eventually degraded in lysosomes, the major catabolic cellular compartment. After endocytosis, the endocytic vesicles carrying the receptors fuse with early endosomes, which then mature to late endosomes containing the material destined for degradation. Mature late endosomes are competent to fuse with lysosomes that contain a variety of hydrolases for the breakdown of all kinds of macromolecules^[100].

There is now solid data that, at the end of their lifetime, GABA_B receptors are endocytosed and degraded in lysosomes. This is evidenced by the intracellular accumulation of internalized GABA_B receptors upon inhibition of lysosomal function^[83,84,101] and the colocalization of intracellular GABA_B receptors with marker proteins for late endosomes and lysosomes^[84,92]. GABA_B receptors are most likely sorted by the ESCRT (endosomal sorting complex required for transport) machinery to lysosomes, because the knockdown of tumor susceptibility gene 101 (TSG101), an integral component of the ESCRT machinery, prevents degradation of the receptors^[101]. Three distinct ESCRT complexes sequentially target mono- and K63-linked polyubiquitinated membrane proteins to late endosomes^[102]. However, it remains to be shown whether GABA_B receptors are ubiquitinated and whether ubiquitination serves as a lysosomal sorting signal.

Another unresolved issue is how the decision is made as to whether a receptor is sorted to the degradation pathway. As discussed above, the vast majority of endocytosed GABA_B receptors recycle back to the plasma membrane and only few are degraded. However, pharmacological inhibition of recycling leads to rapid lysosomal degradation of the receptors (about 50% of the total receptor population within 30 min)^[84]. This indicates that recycling and degradation of GABA_B receptors is tightly controlled, and decreasing the rate of recycling constitutes a mechanism to rapidly reduce the receptor number (discussed below).




Figure 3 Regulation of cell surface GABA_B receptors by trafficking. A: Under normal conditions GABA_B receptors are constitutively internalized and recycled back to the plasma membrane. Only a small fraction of receptors are sorted to lysosomes for degradation; B: Sustained activation of glutamate receptors [primarily 2-amino-3-(5-methyl-3-oxo-1,2-oxazol-4-yl) propanoic acid (AMPA) and N-methyl-D-aspartic acid (NMDA) receptors] and L-type voltage-gated Ca²⁺ channels raises intracellular Ca²⁺ levels. This induces phosphorylation of GABA_{B1} at serine 867 by calmodulin-dependent protein kinase II (CaMK II) and of GABA_{B2} at serine 783 by adenosine monophosphate (AMP) kinase, followed by slow dephosphorylation, by protein phosphatase 2 (PPA2). These events shift the recycling/degradation equilibrium towards degradation so that the majority of GABA_B receptors are no longer recycled, but instead degraded in lysosomes. Since constitutive endocytosis of the receptors remains unaffected, this mechanism results in a rapid down-regulation of GABA_B receptors. AMPK: 5'AMP-dependent protein kinase; GABA: γ -Aminobutyric acid; VSCCs: Voltage-sensitive calcium channels.

Regulation of cell surface GABA_B receptors by glutamatergic excitatory activity

GABA_B receptors control glutamate signaling *via* presynaptic and postsynaptic mechanisms. They are abundantly expressed at glutamatergic synapses^[2-5,103] where they are activated by GABA spillover from adjacent GABAergic terminals and inhibit glutamate release^[8,104-106]. This limits activation of postsynaptically located excitatory glutamate receptors (AMPA/kainate and NMDA receptors). Although GABA_B receptors are also located in close proximity to AMPA and NMDA receptors they do not

appear to directly modulate AMPA and NMDA receptor excitatory postsynaptic currents (EPSCs)^[105]. However, activation of postsynaptic GABA_B receptors seem to limit Ca²⁺-influx through NMDA receptors by inhibition of the cAMP/PKA signaling pathway, which normally enhance NMDA receptor Ca²⁺ conductance^[105].

Besides the prominent regulation of glutamate signaling by GABA_B receptors, there is now evidence that glutamatergic activity, in return, may affect GABA_B receptor expression to attenuate inhibitory control (Figure 3). Application of glutamate to cultured neurons dra-

matically down-regulates cell surface GABA_B receptors and GABA_B receptor-activated currents^[90,94,97,98]. Specific activation of AMPA receptors^[94] or NMDA receptors^[97,98] was sufficient to induce the down-regulation of GABA_B receptors. Interestingly, the kinetics of AMPA-induced down-regulation of GABA_B receptors was significantly slower than that induced by glutamate and was accelerated upon co-activation of group I metabotropic glutamate receptors^[94]. These findings indicate that beside the ionotropic AMPA and NMDA receptors, metabotropic glutamate receptors also contribute to the glutamate-induced down-regulation of GABA_B receptors. The underlying mechanism of this rapid down-regulation of GABA_B receptors is a shift of the recycling/degradation equilibrium towards lysosomal degradation^[94,97]. Glutamate application reduced the rate of GABA_B receptor recycling without altering the rate of their internalization and was fully restored after inhibition of lysosomal degradation. The precise intracellular signaling cascade leading to the glutamate-induced shift in sorting the GABA_B receptors preferentially to the degradation pathway is currently not fully resolved. It is clear that the down-regulation of GABA_B receptors depends on the influx of Ca²⁺^[94,98], which is most likely mediated by L-type voltage-gated Ca²⁺ channels^[94]. Two downstream effector systems were identified to be involved in the down-regulation of GABA_B receptors (Figure 3). One depends on phosphorylation of serine 867 (S867) in GABA_{B1} by Ca²⁺/calmodulin-dependent protein kinase II (CaMK II)^[98]. The other involves phosphorylation of serine 783 (S783) in GABA_{B2} by AMP kinase and subsequent dephosphorylation by protein phosphatase 2A (PP2A)^[97]. Mutational inactivation of each phosphorylation site prevented glutamate-induced down-regulation of GABA_B receptors. However, while the cell surface expression of the receptors containing the mutant GABA_{B1}(S867A) was normal^[98], the mutant GABA_{B2}(S783A) was expressed to a significantly lesser level in the plasma membrane^[97]. This suggests that phosphorylation of S783 in GABA_{B2} is involved in sorting the receptors to the recycling pathway, while phosphorylation of S867 in GABA_{B1} may constitute a direct signal for sorting the receptors to lysosomal degradation. Alternatively, phosphorylation of GABA_{B1}(S867) may be required for dephosphorylation of S783 in GABA_{B2}, for instance by recruiting PPA2 to the receptor. In this respect it would be very interesting to test whether phosphorylation of GABA_{B1}(S867) by CaMK II is required for dephosphorylation of GABA_{B2}(S783) by PPA2.

What is the physiological relevance of this mechanism? Since glutamate-induced down-regulation of GABA_B receptors has so far only been studied in cultured neurons, the role of this process *in vivo* remains to be shown. However, there are physiological, as well as pathological, conditions involving sustained activity of glutamate receptors where this mechanism might be operative. Under pathological conditions associated with excessive activation of glutamate receptors, such as ischemia, down-

regulation of GABA_B receptors results in diminished inhibitory control and may further enhance excitotoxicity and neuronal cell death. This view is supported by an *in vitro* model of ischemia where total GABA_{B2} protein levels were found to be strongly reduced 60 min after the ischemic insult^[107]. Likewise, in an *in vivo* model of hypoxia/ischemia, significantly reduced levels of GABA_B receptors were detected^[108].

Under normal physiological conditions, glutamate-induced down-regulation of GABA_B receptors may contribute to the process of long-term potentiation, which is thought to be the molecular basis for learning and memory formation, as long-term potentiation is associated with sustained activity of glutamate receptors^[109]. In this scenario, enhanced glutamatergic activity would induce the down-regulation of GABA_B receptors and consequently relieve the synapses from inhibition, resulting in a further increase of synaptic excitability.

CONCLUSION

Trafficking events play a pivotal role in the cell surface availability of receptors and largely determine their signaling strength. Currently, we are only beginning to identify and understand the trafficking mechanisms of GABA_B receptors and how cell surface expression of the receptors is regulated. In particular, we almost completely lack knowledge on forward trafficking of GABA_B receptors from the ER *via* the Golgi network to the plasma membrane. In addition, mechanisms on the targeting of the receptors to specific sites in the neuron are unknown. There is an initial indication that GABA_{B1} may be transported independent of GABA_{B2} within the ER into dendrites and are then assembled and exported to the plasma membrane^[110]. This finding implies that heterodimerization of GABA_B receptors is a spatially and temporally controlled mechanism, and would provide an additional level to regulate cell surface expression of the receptors. It is now clear that GABA_B receptors are constitutively endocytosed *via* the clathrin and dynamin-dependent pathway, and are predominantly recycled back to the plasma membrane with only a minor fraction being degraded in lysosomes. The equilibrium of sorting the receptors to the recycling and degradation pathway appears to be controlled by phosphorylation/dephosphorylation events and regulated by changes in neuronal activity associated with increased influx of Ca²⁺. It will be a major future effort to unravel the mechanisms involved in trafficking, sorting and degradation of GABA_B receptors and how they are regulated by physiological and pathological stimuli. It is now well established that receptor trafficking regulates signal transduction and that disturbances in these mechanisms may contribute to disease states^[111]. Since GABA_B receptors have been implicated in a variety of neurological disorders-ranging from epilepsy, addiction, schizophrenia, depression, anxiety to chronic pain-it is likely that altered GABA_B receptor trafficking is involved, at least to some extent, in these diseases. We expect that a deeper knowl-

edge of the trafficking mechanisms of GABA_B receptors under physiological and pathological conditions will provide the basis for the development of novel and highly selective future therapeutic interventions.

REFERENCES

- Ulrich D, Bettler B. GABA(B) receptors: synaptic functions and mechanisms of diversity. *Curr Opin Neurobiol* 2007; **17**: 298-303
- Luján R, Shigemoto R, Kulik A, Juiz JM. Localization of the GABAB receptor 1a/b subunit relative to glutamatergic synapses in the dorsal cochlear nucleus of the rat. *J Comp Neurol* 2004; **475**: 36-46
- Kulik A, Vida I, Luján R, Haas CA, López-Bendito G, Shigemoto R, Frotscher M. Subcellular localization of metabotropic GABA(B) receptor subunits GABA(B1a/b) and GABA(B2) in the rat hippocampus. *J Neurosci* 2003; **23**: 11026-11035
- Lacey CJ, Boyes J, Gerlach O, Chen L, Magill PJ, Bolam JP. GABA(B) receptors at glutamatergic synapses in the rat striatum. *Neuroscience* 2005; **136**: 1083-1095
- Chen L, Boyes J, Yung WH, Bolam JP. Subcellular localization of GABAB receptor subunits in rat globus pallidus. *J Comp Neurol* 2004; **474**: 340-352
- Boyes J, Bolam JP. The subcellular localization of GABA(B) receptor subunits in the rat substantia nigra. *Eur J Neurosci* 2003; **18**: 3279-3293
- Luján R, Shigemoto R. Localization of metabotropic GABA receptor subunits GABAB1 and GABAB2 relative to synaptic sites in the rat developing cerebellum. *Eur J Neurosci* 2006; **23**: 1479-1490
- Scanziani M. GABA spillover activates postsynaptic GABA(B) receptors to control rhythmic hippocampal activity. *Neuron* 2000; **25**: 673-681
- Zilberter Y, Kaiser KM, Sakmann B. Dendritic GABA release depresses excitatory transmission between layer 2/3 pyramidal and bitufted neurons in rat neocortex. *Neuron* 1999; **24**: 979-988
- Angulo MC, Le Meur K, Kozlov AS, Charpak S, Audinat E. GABA, a forgotten gliotransmitter. *Prog Neurobiol* 2008; **86**: 297-303
- Vélez-Fort M, Audinat E, Angulo MC. Central Role of GABA in Neuron-Glia Interactions. *Neuroscientist* 2011; Epub ahead of print
- Laviv T, Riven I, Dolev I, Vertkin I, Balana B, Slesinger PA, Slutsky I. Basal GABA regulates GABA(B)R conformation and release probability at single hippocampal synapses. *Neuron* 2010; **67**: 253-267
- Nishikawa M, Hirouchi M, Kuriyama K. Functional coupling of Gi subtype with GABAB receptor/adenylyl cyclase system: analysis using a reconstituted system with purified GTP-binding protein from bovine cerebral cortex. *Neurochem Int* 1997; **31**: 21-25
- Morishita R, Kato K, Asano T. GABAB receptors couple to G proteins Go, Go* and Gi1 but not to Gi2. *FEBS Lett* 1990; **271**: 231-235
- Menon-Johansson AS, Berrow N, Dolphin AC. G(o) transduces GABAB-receptor modulation of N-type calcium channels in cultured dorsal root ganglion neurons. *Pflugers Arch* 1993; **425**: 335-343
- Odagaki Y, Koyama T. Identification of galpha subtype(s) involved in gamma-aminobutyric acid(B) receptor-mediated high-affinity guanosine triphosphatase activity in rat cerebral cortical membranes. *Neurosci Lett* 2001; **297**: 137-141
- Campbell V, Berrow N, Dolphin AC. GABAB receptor modulation of Ca2+ currents in rat sensory neurones by the G protein G(0): antisense oligonucleotide studies. *J Physiol* 1993; **470**: 1-11
- Mannoury la Cour C, Herbelles C, Pasteau V, de Nanteuil G, Millan MJ. Influence of positive allosteric modulators on GABA(B) receptor coupling in rat brain: a scintillation proximity assay characterisation of G protein subtypes. *J Neurochem* 2008; **105**: 308-323
- Sakaba T, Neher E. Direct modulation of synaptic vesicle priming by GABA(B) receptor activation at a glutamatergic synapse. *Nature* 2003; **424**: 775-778
- Ghorbel MT, Becker KG, Henley JM. Profile of changes in gene expression in cultured hippocampal neurones evoked by the GABAB receptor agonist baclofen. *Physiol Genomics* 2005; **22**: 93-98
- Helm KA, Haberman RP, Dean SL, Hoyt EC, Melcher T, Lund PK, Gallagher M. GABAB receptor antagonist SGS742 improves spatial memory and reduces protein binding to the cAMP response element (CRE) in the hippocampus. *Neuropharmacology* 2005; **48**: 956-964
- Barthel F, Kienlen Campard P, Demeneix BA, Feltz P, Loeffler JP. GABAB receptors negatively regulate transcription in cerebellar granular neurons through cyclic AMP responsive element binding protein-dependent mechanisms. *Neuroscience* 1996; **70**: 417-427
- Guyon A, Leresche N. Modulation by different GABAB receptor types of voltage-activated calcium currents in rat thalamocortical neurones. *J Physiol* 1995; **485** (Pt 1): 29-42
- Mintz IM, Bean BP. GABAB receptor inhibition of P-type Ca2+ channels in central neurons. *Neuron* 1993; **10**: 889-898
- Chen G, van den Pol AN. Presynaptic GABAB autoreceptor modulation of P/Q-type calcium channels and GABA release in rat suprachiasmatic nucleus neurons. *J Neurosci* 1998; **18**: 1913-1922
- Bussi eres N, El Manira A. GABA(B) receptor activation inhibits N- and P/Q-type calcium channels in cultured lamprey sensory neurons. *Brain Res* 1999; **847**: 175-185
- Lambert NA, Wilson WA. High-threshold Ca2+ currents in rat hippocampal interneurons and their selective inhibition by activation of GABA(B) receptors. *J Physiol* 1996; **492** (Pt 1): 115-127
- Bean BP. Neurotransmitter inhibition of neuronal calcium currents by changes in channel voltage dependence. *Nature* 1989; **340**: 153-156
- L uscher C, Jan LY, Stoffel M, Malenka RC, Nicoll RA. G protein-coupled inwardly rectifying K+ channels (GIRKs) mediate postsynaptic but not presynaptic transmitter actions in hippocampal neurons. *Neuron* 1997; **19**: 687-695
- Andrade R, Malenka RC, Nicoll RA. A G protein couples serotonin and GABAB receptors to the same channels in hippocampus. *Science* 1986; **234**: 1261-1265
- Inanobe A, Morishige KI, Takahashi N, Ito H, Yamada M, Takumi T, Nishina H, Takahashi K, Kanaho Y, Katada T. G beta gamma directly binds to the carboxyl terminus of the G protein-gated muscarinic K+ channel, GIRK1. *Biochem Biophys Res Commun* 1995; **212**: 1022-1028
- Huang CL, Slesinger PA, Casey PJ, Jan YN, Jan LY. Evidence that direct binding of G beta gamma to the GIRK1 G protein-gated inwardly rectifying K+ channel is important for channel activation. *Neuron* 1995; **15**: 1133-1143
- Wickman KD, I niguez-Lluhl JA, Davenport PA, Taussig R, Krapivinsky GB, Linder ME, Gilman AG, Clapham DE. Recombinant G-protein beta gamma subunits activate the muscarinic-gated atrial potassium channel. *Nature* 1994; **368**: 255-257
- Reuveny E, Slesinger PA, Inglese J, Morales JM, I niguez-Lluhi JA, Lefkowitz RJ, Bourne HR, Jan YN, Jan LY. Activation of the cloned muscarinic potassium channel by G protein beta gamma subunits. *Nature* 1994; **370**: 143-146
- Saint DA, Thomas T, Gage PW. GABAB agonists modulate a transient potassium current in cultured mammalian hippocampal neurons. *Neurosci Lett* 1990; **118**: 9-13
- Fern andez-Alacid L, Aguado C, Ciriuela F, Mart n R, Col n J, Caba ero MJ, Gassmann M, Watanabe M, Shigemoto R, Wickman K, Bettler B, S nchez-Prieto J, Luj n R. Subcellular

- compartment-specific molecular diversity of pre- and post-synaptic GABA-activated GIRK channels in Purkinje cells. *J Neurochem* 2009; **110**: 1363-1376
- 37 **Ladera C**, del Carmen Godino M, José Cabañero M, Torres M, Watanabe M, Luján R, Sánchez-Prieto J. Pre-synaptic GABA receptors inhibit glutamate release through GIRK channels in rat cerebral cortex. *J Neurochem* 2008; **107**: 1506-1517
 - 38 **Pérez-Garci E**, Gassmann M, Bettler B, Larkum ME. The GABAB1b isoform mediates long-lasting inhibition of dendritic Ca²⁺ spikes in layer 5 somatosensory pyramidal neurons. *Neuron* 2006; **50**: 603-616
 - 39 **Kavalali ET**, Zhuo M, Bito H, Tsien RW. Dendritic Ca²⁺ channels characterized by recordings from isolated hippocampal dendritic segments. *Neuron* 1997; **18**: 651-663
 - 40 **Chalifoux JR**, Carter AG. GABAB receptor modulation of synaptic function. *Curr Opin Neurobiol* 2011; **21**: 339-344
 - 41 **Sabatini BL**, Svoboda K. Analysis of calcium channels in single spines using optical fluctuation analysis. *Nature* 2000; **408**: 589-593
 - 42 **Kulik A**, Vida I, Fukazawa Y, Guetg N, Kasugai Y, Marker CL, Rigato F, Bettler B, Wickman K, Frotscher M, Shigemoto R. Compartment-dependent colocalization of Kir3.2-containing K⁺ channels and GABAB receptors in hippocampal pyramidal cells. *J Neurosci* 2006; **26**: 4289-4297
 - 43 **Park HW**, Jung H, Choi KH, Baik JH, Rhim H. Direct interaction and functional coupling between voltage-gated CaV1.3 Ca²⁺ channel and GABAB receptor subunit 2. *FEBS Lett* 2010; **584**: 3317-3322
 - 44 **Bowery NG**, Hill DR, Hudson AL, Doble A, Middlemiss DN, Shaw J, Turnbull M. (-)Baclofen decreases neurotransmitter release in the mammalian CNS by an action at a novel GABA receptor. *Nature* 1980; **283**: 92-94
 - 45 **Kaupmann K**, Huggel K, Heid J, Flor PJ, Bischoff S, Mickel SJ, McMaster G, Angst C, Bittiger H, Froestl W, Bettler B. Expression cloning of GABA(B) receptors uncovers similarity to metabotropic glutamate receptors. *Nature* 1997; **386**: 239-246
 - 46 **Steiger JL**, Bandyopadhyay S, Farb DH, Russek SJ. cAMP response element-binding protein, activating transcription factor-4, and upstream stimulatory factor differentially control hippocampal GABABR1a and GABABR1b subunit gene expression through alternative promoters. *J Neurosci* 2004; **24**: 6115-6126
 - 47 **Kaupmann K**, Malitschek B, Schuler V, Heid J, Froestl W, Beck P, Mosbacher J, Bischoff S, Kulik A, Shigemoto R, Karschin A, Bettler B. GABA(B)-receptor subtypes assemble into functional heteromeric complexes. *Nature* 1998; **396**: 683-687
 - 48 **Kuner R**, Köhr G, Grünewald S, Eisenhardt G, Bach A, Kornau HC. Role of heteromer formation in GABAB receptor function. *Science* 1999; **283**: 74-77
 - 49 **Jones KA**, Borowsky B, Tamm JA, Craig DA, Durkin MM, Dai M, Yao WJ, Johnson M, Gunwaldsen C, Huang LY, Tang C, Shen Q, Salon JA, Morse K, Laz T, Smith KE, Nagarathnam D, Noble SA, Branchek TA, Gerald C. GABA(B) receptors function as a heteromeric assembly of the subunits GABA(B)R1 and GABA(B)R2. *Nature* 1998; **396**: 674-679
 - 50 **White JH**, Wise A, Main MJ, Green A, Fraser NJ, Disney GH, Barnes AA, Emson P, Foord SM, Marshall FH. Heterodimerization is required for the formation of a functional GABA(B) receptor. *Nature* 1998; **396**: 679-682
 - 51 **Ng GY**, Clark J, Coulombe N, Ethier N, Hebert TE, Sullivan R, Kargman S, Chateaufneuf A, Tsukamoto N, McDonald T, Whiting P, Mezey E, Johnson MP, Liu Q, Kolakowski LF, Evans JF, Bonner TI, O'Neill GP. Identification of a GABAB receptor subunit, gb2, required for functional GABAB receptor activity. *J Biol Chem* 1999; **274**: 7607-7610
 - 52 **Galvez T**, Parmentier ML, Joly C, Malitschek B, Kaupmann K, Kuhn R, Bittiger H, Froestl W, Bettler B, Pin JP. Mutagenesis and modeling of the GABAB receptor extracellular domain support a venus flytrap mechanism for ligand binding. *J Biol Chem* 1999; **274**: 13362-13369
 - 53 **Galvez T**, Prezeau L, Milioti G, Franek M, Joly C, Froestl W, Bettler B, Bertrand HO, Blahos J, Pin JP. Mapping the agonist-binding site of GABAB type 1 subunit sheds light on the activation process of GABAB receptors. *J Biol Chem* 2000; **275**: 41166-41174
 - 54 **Bernard P**, Guedin D, Hibert M. Molecular modeling of the GABA/GABA(B) receptor complex. *J Med Chem* 2001; **44**: 27-35
 - 55 **Liu J**, Maurel D, Etzol S, Brabet I, Ansanay H, Pin JP, Rondard P. Molecular determinants involved in the allosteric control of agonist affinity in the GABAB receptor by the GABAB2 subunit. *J Biol Chem* 2004; **279**: 15824-15830
 - 56 **Galvez T**, Duthey B, Kniazeff J, Blahos J, Rovelli G, Bettler B, Prezeau L, Pin JP. Allosteric interactions between GB1 and GB2 subunits are required for optimal GABA(B) receptor function. *EMBO J* 2001; **20**: 2152-2159
 - 57 **Binet V**, Brajon C, Le Corre L, Acher F, Pin JP, Prezeau L. The heptahelical domain of GABA(B2) is activated directly by CGP7930, a positive allosteric modulator of the GABA(B) receptor. *J Biol Chem* 2004; **279**: 29085-29091
 - 58 **Urwyler S**, Gjoni T, Koljatic J, Dupuis DS. Mechanisms of allosteric modulation at GABAB receptors by CGP7930 and GS39783: effects on affinities and efficacies of orthosteric ligands with distinct intrinsic properties. *Neuropharmacology* 2005; **48**: 343-353
 - 59 **Duthey B**, Caudron S, Perroy J, Bettler B, Fagni L, Pin JP, Prezeau L. A single subunit (GB2) is required for G-protein activation by the heterodimeric GABA(B) receptor. *J Biol Chem* 2002; **277**: 3236-3241
 - 60 **Havlickova M**, Prezeau L, Duthey B, Bettler B, Pin JP, Blahos J. The intracellular loops of the GB2 subunit are crucial for G-protein coupling of the heteromeric gamma-aminobutyrate B receptor. *Mol Pharmacol* 2002; **62**: 343-350
 - 61 **Robbins MJ**, Calver AR, Filippov AK, Hirst WD, Russell RB, Wood MD, Nasir S, Couve A, Brown DA, Moss SJ, Pangalos MN. GABA(B2) is essential for g-protein coupling of the GABA(B) receptor heterodimer. *J Neurosci* 2001; **21**: 8043-8052
 - 62 **Margeta-Mitrovic M**, Jan YN, Jan LY. Ligand-induced signal transduction within heterodimeric GABA(B) receptor. *Proc Natl Acad Sci USA* 2001; **98**: 14643-14648
 - 63 **Margeta-Mitrovic M**, Jan YN, Jan LY. Function of GB1 and GB2 subunits in G protein coupling of GABA(B) receptors. *Proc Natl Acad Sci USA* 2001; **98**: 14649-14654
 - 64 **Margeta-Mitrovic M**, Jan YN, Jan LY. A trafficking checkpoint controls GABA(B) receptor heterodimerization. *Neuron* 2000; **27**: 97-106
 - 65 **Pagano A**, Rovelli G, Mosbacher J, Lohmann T, Duthey B, Stauffer D, Ristig D, Schuler V, Meigel I, Lampert C, Stein T, Prezeau L, Blahos J, Pin J, Froestl W, Kuhn R, Heid J, Kaupmann K, Bettler B. C-terminal interaction is essential for surface trafficking but not for heteromeric assembly of GABA(b) receptors. *J Neurosci* 2001; **21**: 1189-1202
 - 66 **Calver AR**, Robbins MJ, Cosio C, Rice SQ, Babbs AJ, Hirst WD, Boyfield I, Wood MD, Russell RB, Price GW, Couve A, Moss SJ, Pangalos MN. The C-terminal domains of the GABA(b) receptor subunits mediate intracellular trafficking but are not required for receptor signaling. *J Neurosci* 2001; **21**: 1203-1210
 - 67 **Couve A**, Filippov AK, Connolly CN, Bettler B, Brown DA, Moss SJ. Intracellular retention of recombinant GABAB receptors. *J Biol Chem* 1998; **273**: 26361-26367
 - 68 **Gassmann M**, Haller C, Stoll Y, Abdel Aziz S, Biermann B, Mosbacher J, Kaupmann K, Bettler B. The RXR-type endoplasmic reticulum-retention/retrieval signal of GABAB1 requires distant spacing from the membrane to function. *Mol Pharmacol* 2005; **68**: 137-144
 - 69 **Gainetdinov RR**, Premont RT, Bohn LM, Lefkowitz RJ,

- Caron MG. Desensitization of G protein-coupled receptors and neuronal functions. *Annu Rev Neurosci* 2004; **27**: 107-144
- 70 **Ferguson SS**. Evolving concepts in G protein-coupled receptor endocytosis: the role in receptor desensitization and signaling. *Pharmacol Rev* 2001; **53**: 1-24
- 71 **Couve A**, Thomas P, Calver AR, Hirst WD, Pangalos MN, Walsh FS, Smart TG, Moss SJ. Cyclic AMP-dependent protein kinase phosphorylation facilitates GABA(B) receptor-effector coupling. *Nat Neurosci* 2002; **5**: 415-424
- 72 **González-Maeso J**, Wise A, Green A, Koenig JA. Agonist-induced desensitization and endocytosis of heterodimeric GABAB receptors in CHO-K1 cells. *Eur J Pharmacol* 2003; **481**: 15-23
- 73 **Tosetti P**, Bakels R, Colin-Le Brun I, Ferrand N, Gaiarsa JL, Caillard O. Acute desensitization of presynaptic GABAB-mediated inhibition and induction of epileptiform discharges in the neonatal rat hippocampus. *Eur J Neurosci* 2004; **19**: 3227-3234
- 74 **Pontier SM**, Lahaie N, Ginham R, St-Gelais F, Bonin H, Bell DJ, Flynn H, Trudeau LE, McIlhinney J, White JH, Bouvier M. Coordinated action of NSF and PKC regulates GABAB receptor signaling efficacy. *EMBO J* 2006; **25**: 2698-2709
- 75 **Kanaide M**, Uezono Y, Matsumoto M, Hojo M, Ando Y, Sudo Y, Sumikawa K, Taniyama K. Desensitization of GABA(B) receptor signaling by formation of protein complexes of GABA(B2) subunit with GRK4 or GRK5. *J Cell Physiol* 2007; **210**: 237-245
- 76 **Virlon B**, Firsov D, Cheval L, Reiter E, Troispoux C, Guillou F, Elalouf JM. Rat G protein-coupled receptor kinase GRK4: identification, functional expression, and differential tissue distribution of two splice variants. *Endocrinology* 1998; **139**: 2784-2795
- 77 **Sallese M**, Salvatore L, D'Urbano E, Sala G, Storto M, Launey T, Nicoletti F, Knöpfel T, De Blasi A. The G-protein-coupled receptor kinase GRK4 mediates homologous desensitization of metabotropic glutamate receptor 1. *FASEB J* 2000; **14**: 2569-2580
- 78 **Perroy J**, Adam L, Qanbar R, Chénier S, Bouvier M. Phosphorylation-independent desensitization of GABA(B) receptor by GRK4. *EMBO J* 2003; **22**: 3816-3824
- 79 **Ando Y**, Hojo M, Kanaide M, Takada M, Sudo Y, Shiraishi S, Sumikawa K, Uezono Y. S(+)-ketamine suppresses desensitization of γ -aminobutyric acid type B receptor-mediated signaling by inhibition of the interaction of γ -aminobutyric acid type B receptors with G protein-coupled receptor kinase 4 or 5. *Anesthesiology* 2011; **114**: 401-411
- 80 **Schwenk J**, Metz M, Zolles G, Turecek R, Fritzius T, Bildl W, Tarusawa E, Kulik A, Unger A, Ivankova K, Seddik R, Tiao JY, Rajalu M, Trojanova J, Rohde V, Gassmann M, Schulte U, Fakler B, Bettler B. Native GABA(B) receptors are heteromultimers with a family of auxiliary subunits. *Nature* 2010; **465**: 231-235
- 81 **Cruz HG**, Ivanova T, Lunn ML, Stoffel M, Slesinger PA, Lüscher C. Bi-directional effects of GABA(B) receptor agonists on the mesolimbic dopamine system. *Nat Neurosci* 2004; **7**: 153-159
- 82 **Fairfax BP**, Pitcher JA, Scott MG, Calver AR, Pangalos MN, Moss SJ, Couve A. Phosphorylation and chronic agonist treatment atypically modulate GABAB receptor cell surface stability. *J Biol Chem* 2004; **279**: 12565-12573
- 83 **Grampp T**, Sauter K, Markovic B, Benke D. Gamma-aminobutyric acid type B receptors are constitutively internalized via the clathrin-dependent pathway and targeted to lysosomes for degradation. *J Biol Chem* 2007; **282**: 24157-24165
- 84 **Grampp T**, Notz V, Broll I, Fischer N, Benke D. Constitutive, agonist-accelerated, recycling and lysosomal degradation of GABA(B) receptors in cortical neurons. *Mol Cell Neurosci* 2008; **39**: 628-637
- 85 **Mutneja M**, Berton F, Suen KF, Lüscher C, Slesinger PA. Endogenous RGS proteins enhance acute desensitization of GABA(B) receptor-activated GIRK currents in HEK-293T cells. *Pflugers Arch* 2005; **450**: 61-73
- 86 **Kuramoto N**, Wilkins ME, Fairfax BP, Revilla-Sanchez R, Terunuma M, Tamaki K, Iemata M, Warren N, Couve A, Calver A, Horvath Z, Freeman K, Carling D, Huang L, Gonzales C, Cooper E, Smart TG, Pangalos MN, Moss SJ. Phospho-dependent functional modulation of GABA(B) receptors by the metabolic sensor AMP-dependent protein kinase. *Neuron* 2007; **53**: 233-247
- 87 **Carling D**. AMP-activated protein kinase: balancing the scales. *Biochimie* 2005; **87**: 87-91
- 88 **Kahn BB**, Alquier T, Carling D, Hardie DG. AMP-activated protein kinase: ancient energy gauge provides clues to modern understanding of metabolism. *Cell Metab* 2005; **1**: 15-25
- 89 **Brock C**, Boudier L, Maurel D, Blahos J, Pin JP. Assembly-dependent surface targeting of the heterodimeric GABAB Receptor is controlled by COPI but not 14-3-3. *Mol Biol Cell* 2005; **16**: 5572-5578
- 90 **Vargas KJ**, Terunuma M, Tello JA, Pangalos MN, Moss SJ, Couve A. The availability of surface GABA B receptors is independent of gamma-aminobutyric acid but controlled by glutamate in central neurons. *J Biol Chem* 2008; **283**: 24641-24648
- 91 **Pooler AM**, Gray AG, McIlhinney RA. Identification of a novel region of the GABA(B2) C-terminus that regulates surface expression and neuronal targeting of the GABA(B) receptor. *Eur J Neurosci* 2009; **29**: 869-878
- 92 **Hannan S**, Wilkins ME, Dehghani-Tafti E, Thomas P, Baddeley SM, Smart TG. GABAB receptor internalisation is regulated by the R2 subunit. *J Biol Chem* 2011; Epub ahead of print
- 93 **Wilkins ME**, Li X, Smart TG. Tracking cell surface GABAB receptors using an alpha-bungarotoxin tag. *J Biol Chem* 2008; **283**: 34745-34752
- 94 **Maier PJ**, Marin I, Grampp T, Sommer A, Benke D. Sustained glutamate receptor activation down-regulates GABAB receptors by shifting the balance from recycling to lysosomal degradation. *J Biol Chem* 2010; **285**: 35606-35614
- 95 **Laffray S**, Tan K, Dulluc J, Bouali-Benazzouz R, Calver AR, Nagy F, Landry M. Dissociation and trafficking of rat GABAB receptor heterodimer upon chronic capsaicin stimulation. *Eur J Neurosci* 2007; **25**: 1402-1416
- 96 **Ungewickell EJ**, Hinrichsen L. Endocytosis: clathrin-mediated membrane budding. *Curr Opin Cell Biol* 2007; **19**: 417-425
- 97 **Terunuma M**, Vargas KJ, Wilkins ME, Ramírez OA, Jaureguierry-Bravo M, Pangalos MN, Smart TG, Moss SJ, Couve A. Prolonged activation of NMDA receptors promotes dephosphorylation and alters postendocytic sorting of GABAB receptors. *Proc Natl Acad Sci USA* 2010; **107**: 13918-13923
- 98 **Guett N**, Abdel Aziz S, Holbro N, Turecek R, Rose T, Seddik R, Gassmann M, Moes S, Jenoe P, Oertner TG, Casanova E, Bettler B. NMDA receptor-dependent GABAB receptor internalization via CaMKII phosphorylation of serine 867 in GABAB1. *Proc Natl Acad Sci USA* 2010; **107**: 13924-13929
- 99 **Park M**, Penick EC, Edwards JG, Kauer JA, Ehlers MD. Recycling endosomes supply AMPA receptors for LTP. *Science* 2004; **305**: 1972-1975
- 100 **Saftig P**, Klumperman J. Lysosome biogenesis and lysosomal membrane proteins: trafficking meets function. *Nat Rev Mol Cell Biol* 2009; **10**: 623-635
- 101 **Kantamneni S**, Holman D, Wilkinson KA, Corrêa SA, Feligioni M, Ogden S, Fraser W, Nishimune A, Henley JM. GISP binding to TSG101 increases GABA receptor stability by down-regulating ESCRT-mediated lysosomal degradation. *J Neurochem* 2008; **107**: 86-95
- 102 **Raiborg C**, Stenmark H. The ESCRT machinery in endosomal sorting of ubiquitylated membrane proteins. *Nature* 2009; **458**: 445-452
- 103 **Vigot R**, Barbieri S, Bräuner-Osborne H, Turecek R, Shi-

- gemoto R, Zhang YP, Luján R, Jacobson LH, Biermann B, Fritschy JM, Vacher CM, Müller M, Sansig G, Guetg N, Cryan JF, Kaupmann K, Gassmann M, Oertner TG, Bettler B. Differential compartmentalization and distinct functions of GABAB receptor variants. *Neuron* 2006; **50**: 589-601
- 104 **Vogt KE**, Nicoll RA. Glutamate and gamma-aminobutyric acid mediate a heterosynaptic depression at mossy fiber synapses in the hippocampus. *Proc Natl Acad Sci USA* 1999; **96**: 1118-1122
- 105 **Chalifoux JR**, Carter AG. GABAB receptors modulate NMDA receptor calcium signals in dendritic spines. *Neuron* 2010; **66**: 101-113
- 106 **Pan BX**, Dong Y, Ito W, Yanagawa Y, Shigemoto R, Morozov A. Selective gating of glutamatergic inputs to excitatory neurons of amygdala by presynaptic GABA_B receptor. *Neuron* 2009; **61**: 917-929
- 107 **Cimarosti H**, Kantamneni S, Henley JM. Ischaemia differentially regulates GABA(B) receptor subunits in organotypic hippocampal slice cultures. *Neuropharmacology* 2009; **56**: 1088-1096
- 108 **Anju TR**, Abraham PM, Antony S, Paulose CS. Alterations in cortical GABAB receptors in neonatal rats exposed to hypoxic stress: role of glucose, oxygen, and epinephrine resuscitation. *Mol Cell Biochem* 2010; **343**: 1-11
- 109 **Blundon JA**, Zakharenko SS. Dissecting the components of long-term potentiation. *Neuroscientist* 2008; **14**: 598-608
- 110 **Ramírez OA**, Vidal RL, Tello JA, Vargas KJ, Kindler S, Härtel S, Couve A. Dendritic assembly of heteromeric gamma-aminobutyric acid type B receptor subunits in hippocampal neurons. *J Biol Chem* 2009; **284**: 13077-13085
- 111 **Scita G**, Di Fiore PP. The endocytic matrix. *Nature* 2010; **463**: 464-473

S- Editor Cheng JX L- Editor Rutherford A E- Editor Zhang DN